

Review

Estrogen receptor transcription and transactivation Estrogen receptor knockout mice: what their phenotypes reveal about mechanisms of estrogen action

Sylvia Curtis Hewitt, John F Couse and Kenneth S Korach

National Institutes of Health, Research Triangle Park, North Carolina, USA

Received: 14 February 2000
Revisions requested: 1 March 2000
Revisions received: 9 March 2000
Accepted: 26 May 2000
Published: 12 July 2000

Breast Cancer Res 2000, **2**:345–352

Current Science Ltd (Print ISSN 1465-5411; Online ISSN 1465-542X)

Abstract

Natural, synthetic and environmental estrogens have numerous effects on the development and physiology of mammals. Estrogen is primarily known for its role in the development and functioning of the female reproductive system. However, roles for estrogen in male fertility, bone, the circulatory system and immune system have been established by clinical observations regarding sex differences in pathologies, as well as observations following menopause or castration. The primary mechanism of estrogen action is via binding and modulation of activity of the estrogen receptors (ERs), which are ligand-dependent nuclear transcription factors. ERs are found in highest levels in female tissues critical to reproduction, including the ovaries, uterus, cervix, mammary glands and pituitary gland. Since other affected tissues have extremely low levels of ER, indirect effects of estrogen, for example induction of pituitary hormones that affect the bone, have been proposed. The development of transgenic mouse models that lack either estrogen or ER have proven to be valuable tools in defining the mechanisms by which estrogen exerts its effects in various systems. The aim of this article is to review the mouse models with disrupted estrogen signaling and describe the associated phenotypes.


Keywords: estrogen receptor, estrogen receptor knockout, transgenic

Reproductive phenotypes of estrogen receptor knockout models

Estrogen has many roles in reproduction, and the generation of the estrogen receptor (ER) α and ER β knockout (α ERKO and β ERKO) mice has further illustrated its roles and mechanisms. Interestingly, both sexes of the α ERKO mice are infertile, whereas only the β ERKO female has shown impaired fertility. In the male α ERKO mice, infertility is due to deficits at several points in the reproductive

process, including severe reduction in sperm numbers and lack of sperm function, as well as abnormal sexual behavior. The seminiferous tubules of the α ERKO testes show progressive dilation that is accompanied by degeneration of the seminiferous epithelium (Fig. 1) [1*,2*]. Transplanted α ERKO sperm was functional when developed in normal host testes [3**]. In contrast, the testes of the β ERKO mice appear normal (Fig. 1), and produce sufficient and functional sperm to allow fertility, resulting in

Figure 1




Pathology of adult α ERKO and β ERKO testes. Sections from wild-type and ER-disrupted testes were stained with hematoxylin and eosin for comparison of their pathology. The wild-type and β ERKO testes are indistinguishable, while the α ERKO testis shows degeneration of the testicular structures.

production of offspring in mice examined to date. Therefore, ER α appears to be more critical than ER β in mediation of the estrogen actions necessary for maintenance of healthy testicular structures and the somatic cell function required for successful sperm maturation.

Normally, the female rodent reproductive tract grows and matures in response to cycling ovarian hormones, including estradiol. The growth and maturation of the epithelial portion and the preparation of the stromal layer is thought to be important for successful implantation and pregnancy to occur. The infertility of the female α ERKO mouse is due in part to the insensitivity of the uterus to the mitogenic and differentiative actions of estrogen [4**,5**] (Fig. 2). Microscopic evaluation of the α ERKO uterine tissue indicates that all expected tissues are present but appear immature, as illustrated by a reduced number of glands in the endometrium (Fig. 3). ER α is thus not necessary for development of the uterus, but is necessary for complete maturation and function of the tissue. In contrast, the wild-type and β ERKO uteri are indistinguishable, and show normal organization and development of the stromal, myometrial and epithelial layers (Fig. 3), as well as glandular structures. ER β is thus apparently not required for normal development of the female reproductive tract. Furthermore, when challenged with estrogenic compounds, the wild-type and

Figure 2



Gross morphology of adult ERKO female reproductive tracts. Reproductive tracts dissected from wild-type and β ERKO animals are normal, while the α ERKO uterus is immature and the ovaries are enlarged and dark-colored due to hemorrhagic cysts.


β ERKO uteri respond comparably with increased weight and epithelial development, while the α ERKO uterus is nonresponsive. The uterus of the β ERKO mouse is fully functional, as pregnancies are successfully carried to term and delivered. The vaginal tissue of the β ERKO female is also similar in the wild type, with indications of cornification, an estrogen response, whereas the vaginal tissue of the α ERKO female remains immature and is not cornified.

Ovary and ovarian hormones

Numerous intraovarian effects of locally synthesized estrogens have been described and postulated to be essential to ovarian function, including modifications in ER levels [6], DNA synthesis and cell proliferation [7–11], intercellular gap junctions [12], and follicular atresia [13]. Estradiol is also known to augment the actions of follicle-stimulating hormone on granulosa cells, resulting in the maintenance of follicle-stimulating hormone receptor levels [14,15] and the acquisition of luteinizing hormone (LH) receptor [16,17], an event critical to successful ovulation. The intraovarian actions of estradiol act to ultimately enhance follicular responsiveness to gonadotropins, and thereby result in increased aromatase activity and further estrogen synthesis [17,18]. Therefore, given the many speculated intraovarian actions of 17 β -estradiol, disruption of the respective ER genes may be expected to result in distinct ovarian phenotypes.

The ovarian phenotypes are a major component of the infertility in the α ERKO mice and the subfertility in the β ERKO mice. The α ERKO female does not ovulate, while the β ERKO female is subfertile with reduced litter numbers and smaller litter sizes compared with wild-type littermates. Interestingly, although both ER α and ER β are detected in the ovary, their localization differs with ER β in the granulosa cells and ER α in the theca and interstitial

Figure 3



Uterine and vaginal histology of adult ERKO mice. Histological analysis of the uterus (top panels) and vagina (bottom panels) shows the β ERKO tissue is indistinguishable from the wild type, showing the normal organization of the uterine tissue into the epithelial (Ep), stromal (St) and myometrial (My) compartments. In contrast, the α ERKO uterine tissue is composed of all three layers, yet they are all immature and hypoplastic. Note also the glands are fewer in number. The vagina of the wild-type and β ERKO mice is identical and cornified, indicating response to estrogen, while the α ERKO vagina shows no cornification. (Reproduced with permission from Couse and Korach [5**].)

cells of the ovary [19*] (Fig. 4). The differential expression of these receptors makes compensatory activity of one receptor in the absence of the other unlikely.

The hallmark phenotype of the α ERKO female is the enlarged hemorrhagic cystic ovary (Fig. 5), although the prepubertal α ERKO ovary looks similar to its wild-type littermate. This phenotype begins to develop progressively as the animal matures and is apparently due to a lack of estradiol feedback inhibition in the pituitary, which results in chronically elevated LH and subsequent hyperstimulation of the ovary (Table 1). This indicates that $ER\alpha$ is responsible for mediating the LH feedback inhibition in the hypothalamic–pituitary axis. The constant LH stimulation in the α ERKO mice results in an abnormal endocrine environment in the α ERKO female, with elevated estradiol and testosterone, and chronic preovulatory basal progesterone levels (Table 1).

The β ERKO ovaries produce normal serum levels of estradiol and testosterone, and the circulating serum gonadotropin levels are also normal (Table 2). However, the β ERKO ovaries function suboptimally, as illustrated by the appearance of numerous unruptured follicles following superovulation. Attempts to superovulate the β ERKO

female results in some ovulation, but the number of oocytes released is reduced compared with wild-type females (Table 3). A role for $ER\beta$ in ovulation is thus indicated, but the mechanism is still being defined.


Mammary gland

The mammary gland develops and functions in response to ovarian hormones, most notably estrogen and progesterone [20]. The female mammary gland is immature at birth and consists of a mainly stromal tissue, with only a rudimentary epithelial duct structure emanating from the nipple. The ducts elongate in response to ovarian and pituitary hormones, eventually filling the stromal tissue with a branched tree-like structure. Lobular alveolar buds develop along the length of these ducts during pregnancy and differentiate into secretory lactational structures.

Transgenic knockout models have been very informative in understanding the roles of estrogen and progesterone in mammary gland development. The role of progesterone is indicated by the lack of development of alveolar buds in the progesterone receptor knockout mouse [21**]. Similarly, the role for $ER\alpha$ is illustrated by the lack of pubertal growth of the epithelial ductal rudiment in the α ERKO female despite elevated circulating serum estradiol levels [22,23]

Figure 4


Wild-type Mouse Ovary: Estrogen Receptor Expression



Localization of ER α and ER β protein in the ovary. Serial sections from mouse ovary were stained with antibody to ER α (top panel) or ER β (bottom panel). Note that ER β immunoreactivity is confined to the granulosa cells of the follicle, while ER α immunoreactivity is localized in the thecal and interstitial cells of the ovary.

(Table 1 and Fig. 6). Transplantation of a wild-type pituitary under the kidney capsule of an α ERKO female resulted in growth and development of the α ERKO mammary epithelium. Removal of ovarian steroids by castration prevented the growth and development of mammary epithelium following pituitary transplant (Bocchinfuso *et al*, submitted).

Figure 5



Ovarian pathology of the ERKO mice. Histological analysis of the wild-type ovary shows normal follicular development and indications of ovulation. The α ERKO ovary shows large cystic structures and arrested follicle development with no indication of ovulation, while the β ERKO ovary shows development of follicles is occurring but with little indication of successful ovulation.

Administration of progesterone together with estradiol to an ovexed α ERKO mouse resulted in alveolar development and some ductal elongation (Bocchinfuso *et al*, submitted). These observations indicate that the α ERKO pituitary in the absence of ovarian hormones is incapable of providing a hormonal environment necessary for mammary development. In addition, the α ERKO mammary gland epithelium is capable of responding with growth when a normally functioning pituitary is present to provide the necessary signals to the ovary and the mammary gland. Finally, it is apparent that the secretion of ovarian hormones is a required element of ductal elongation in the α ERKO mouse. The β ERKO mammary gland is similar to wild-type littermates [5**] (Fig. 6) and is functional because β ERKO mothers are able to nurse their litters. Our observations suggest that ER β is not required for the structural and functional development of the mammary gland.

Table 1

Serum hormone levels in adult wild-type and α ERKO mice

Hormone	Female		Male	
	Wild type	α ERKO	Wild type	α ERKO
Gonadal steroids				
Estradiol (pg/ml)†	29.5 ± 2.5	84.3 ± 12.5*	11.8 ± 3.4	12.9 ± 3.4
Progesterone (ng/ml)†	2.3 ± 0.6	4.0 ± 1.1	0.5 ± 0.3	0.3 ± 0.1
Testosterone (ng/ml)	0.4 ± 0.4	3.2 ± 0.6	9.3 ± 4.0	16.0 ± 2.3
Anterior pituitary				
LH (ng/ml)	0.3 ± 0.04	1.7 ± 0.3*	2.4 ± 1.2	3.7 ± 0.7
FSH (ng/ml)	4.9 ± 0.6	5.4 ± 0.7	26.0 ± 1.4	30.0 ± 1.1
Prolactin (ng/ml)	18.8 ± 10.7	3.5 ± 1.3	ND	ND

Data presented as mean ± standard error of the mean. ERKO, Estrogen receptor knockout; FSH, follicle stimulating hormone; LH, luteinizing hormone; ND, not determined. †These values in the female are different than those reported by Couse *et al* [43*], which were carried out on pooled sera. The values above are the means from assays on individual samples and therefore are more likely to reflect the true levels in the two genotypes. * *t*-test, wild type versus ERKO, *P* < 0.001. Reproduced with permission from Couse and Korach [5**].

Table 2

Phenotypes of α ERKO and β ERKO mice

Tissue	Observation	
	α ERKO	β ERKO
Testis	Progressive dilation and degradation of tubules, low sperm count, nonfunctional sperm	Normal structure, normal sperm count and fertility
Uterus	Immature, unresponsive to estradiol	Normal development and response to estrogen
Ovary	Enlarged, hemorrhagic cysts, follicles arrested at preantral stage, no corpora lutea, no ovulation, elevated serum estrogen and T levels	Subfertile, infrequent and inefficient ovulation; normal estrogen and T
Mammary, female	Ducts do not develop beyond epithelial rudiment at nipple, no alveolar development	Normal, fully functional. Able to nurse offspring
Pituitary	FSH β , LH β , α GSU, mRNAs all elevated, prolactin mRNA reduced	Normal serum gonadotropin levels
Cardiovascular (male)	Lower basal nitric oxide, estrogen protection in vascular injury not lost. Increase in calcium channels, delayed cardiac depolarization	?
Bone	Shorter; female, smaller diameter; male, lower density	Normal
Brain	Male, no intromission, ejaculation decreased aggression; female, no receptive behaviors	Normal sexual behavior

ERKO, Estrogen receptor knockout; FSH, follicle-stimulating hormone; α GSU, gonadotrophin subunit alpha; LH, luteinizing hormone; T, testosterone.

Table 3

Fertility and superovulation data in the β ERKO female mice

Genotype	Continuous mating results			Superovulation results		
	<i>n</i>	Litters/female	Pups/litter	<i>n</i>	Mean	Range
Wild type	6	2.8 \pm 0.4	8.8 \pm 2.5	10	33.7 \pm 4.8	9–57
Heterozygous	ND	ND	ND	11	52.5 \pm 5.7*	20–77
β ERKO	11	1.7 \pm 1.0*	3.1 \pm 1.8**	11	6.0 \pm 1.5*	0–13

Data presented as mean \pm standard error of the mean. ERKO, Estrogen receptor knockout; ND, not determined. Reproduced with permission from Couse and Korach [5**]. * $P < 0.05$ and ** $P < 0.001$ Student's two tailed *t*-test versus wild type.


Skeletal and cardiovascular tissue

Clinical as well as experimental data imply a role for estrogen as well as other endocrine factors in maintaining bone mass. For example, menopause or castration increases the rate of osteoporosis progression in women [24]. The ERKO model is a very useful tool for elucidating the mechanism of estrogen effects in bone and whether these effects are mediated, directly or indirectly, through ERs. ER α is present at very low levels in the bone cells. In the α ERKO female, the bone is normal in terms of density, but it is significantly shorter and smaller in diameter in both sexes of the α ERKO mice. This indicates the possibility of a role for estrogen in bone lengthening [5**]. A recent report has indicated α ERKO mice have insulin-like growth factor-1 levels 30% lower than normal mice [25], which might account for the

shorter bone. However, the altered hormonal environment (elevated estradiol and testosterone; see Table 1) as well as the increased body weight of the α ERKO mouse should also be considered when interpreting this phenotype. The α ERKO males do show a decrease in bone mineral density, a phenotype that is similar to one clinical case of estrogen insensitivity in a man [26]. Our current evaluation of the bones of the β ERKO mouse shows no apparent difference from the wild type (unpublished data), indicating the lack of a major role for ER β in bone physiology.

A role for estrogen in the cardiovascular system has also been implied by clinical observations correlating hormonal status with cardiovascular disease risk factors [27]. Additionally, ER has been detected in the vasculature [28]. In

Figure 6



Mammmary gland whole mounts from ERKO mice. An adult β ERKO mouse displays a fully developed ductal network similar to the wild type. In contrast, the α ERKO mouse has only a rudimentary underdeveloped epithelial duct (arrow).

the α ERKO male heart, an increased number of calcium channels [29^{*}] and an associated delay in cardiac depolar-

ization have been observed. In the α ERKO vasculature, lower basal nitric oxide activity is detected [30] and, interestingly, the ability of estrogen to play a protective role in a vascular injury model is not lost in the α ERKO [31^{*}] or the β ERKO [32^{*}] mice. These observations suggest either that each receptor can compensate for the loss of the other or that another mechanism results in protection.

Progesterone action in the α ERKO female

The activities of estrogen and progesterone are interdependent in the female reproductive cycle [33]. The level of estrogen in the serum peaks just prior to ovulation. This estrogen 'spike' acts upon the uterus, initiating growth and maturation in preparation for pregnancy. Following ovulation, the level of progesterone rises and prepares the uterine stroma for implantation and decidualization, which is a transformation of the stromal cells in response to apposition of the conceptus. Both hormones have been shown to be essential for successful ovulation and pregnancy to occur. The actions of progesterone, like those of estrogen, are mediated through a specific nuclear transcription factor, the progesterone receptor (PR). Since the level of PR is regulated, in part, by estrogen, the activity of progesterone in α ERKO tissues has been studied to determine whether disruption of α ER signaling alters PR signaling. These observations have been reported in detail [34^{**}], but will be summarized here. PR is present at a basal level in the α ERKO uterus at about 60% of the wild-type level, and its mRNA is not increased by estrogen treatment. However, this basal level of PR is sufficient to induce the mRNA of the progesterone-responsive calcitonin and amphiregulin genes. The α ERKO uterus can also be induced to undergo a decidual reaction in response to progesterone and physical trauma, indicating that PR is fully functional both biochemically and physiologically, and is not dependent on ER α for expression or function. Interestingly, the decidual reaction is estrogen dependent in the wild-type mice, but not in the α ERKO mice.

The mammary gland also responds to progesterone, with development of lobular alveolar structures in the epithelium in preparation for lactation. Although the mammary epithelium of the α ERKO mouse is underdeveloped, it can be stimulated with progesterone to develop lobular alveolar structures, suggesting that although the PR is regulated by estrogen, the basal level of PR present in the α ERKO mammary gland is sufficient to mediate progesterone action. The necessity for estrogen induction of PR for progesterone responsiveness is brought into question by these observations. As reported previously [34^{**}], it is possible that the need for estrogen priming is lost in the α ERKO mice because of developmental differences. Although these progesterone responses have not been characterized in the β ERKO mice, females can carry pregnancies to term and lactate normally, implying that the ability of the uterus and mammary gland to respond to progesterone is normal.

Ligand-independent signaling

Numerous studies have shown that nuclear steroid receptors can be activated by membrane receptor pathways to mediate genomic and physiologic responses in the absence of steroid ligand (reviewed in [35**]). This mechanism is initiated when growth factors bind to and activate their membrane receptor, resulting in activation of a phosphorylation cascade, culminating in modulation of nuclear receptor activity, most likely by phosphorylation of the nuclear receptor protein. These studies are summarized and described in detail elsewhere [36*]. In the case of ER α , we have reported activation by either epidermal growth factor (EGF) or insulin-like growth factor-1 [36*,37,38*,39–41]. The α ERKO mouse has allowed us to show that estrogen-like effects of EGF require the ER α *in vivo*, as EGF induced PR mRNA and thymidine incorporation only in uteri from wild-type mice, indicating that the α ERKO mouse was totally unresponsive. Further biochemical studies indicate that the EGF receptor is present and the pathway is functional in the α ERKO mouse, as EGF induced cFOS, an EGF-responsive gene [36*].

Summary

The phenotypes observed in the α ERKO and β ERKO mice illustrate the roles of ER α and ER β in both reproductive and nonreproductive tissues. Although some phenotypes are downstream results of the lack of estrogen signaling (ie chronic LH stimulation of the α ERKO ovary due to loss of estrogen-mediated negative feedback regulation of LH in the pituitary), we have learned a great deal about the roles of ER α and ER β in development and physiology. Continued evaluation and characterization of the phenotypes of both classical estrogen target tissues, as well as other organ systems, should help to uncover previously unconsidered links of estrogen and other signaling pathways. Finally, our recent generation of mice lacking both ERs ($\alpha\beta$ ERKO) will be useful in detecting any interaction between these receptors or compensatory mechanisms of one receptor in the absence of the other, as well as uncovering nonreceptor-mediated estrogen actions [42**].

References

Articles of particular interest have been highlighted as:

- of special interest
- of outstanding interest

1. Eddy EM, Washburn TF, Bunch DO, Goulding EH, Gladen BC, Lubahn DB, Korach KS: **Targeted disruption of the estrogen receptor gene in male mice causes alteration of spermatogenesis and infertility.** *Endocrinology* 1996, **137**:4796–4805.

This study describes the phenotype seen in the α ERKO males, which was unexpected because a role for estrogen in male reproduction was not predicted.

2. Hess RA, Bunick D, Lee KH, Bahr J, Taylor JA, Korach KS, Lubahn DB: **A role for oestrogens in the male reproductive system.** *Nature* 1997, **390**:509–512.

This study describes the mechanism leading to infertility in the α ERKO male – the luminal fluid is not reabsorbed, further defining the role of estrogen in male reproduction.

3. Mahato D, Goulding EH, Korach KS, Eddy EM: **Spermatogenic cells do not require estrogen receptor alpha for development or function.** *Endocrinology* 2000, **141**:1273–1276.

A sophisticated study in which α ERKO germ cells are grown in the testis of a normal donor, resulting in fertile sperm. This indicates that the role of estrogen is not in production of fertile sperm *per se*, but in providing the proper structures and processes for maturation to occur successfully.

4. Lubahn DB, Moyer JS, Golding TS, Couse JF, Korach KS, Smithies O: **Alteration of reproductive function but not prenatal sexual development after insertional disruption of the mouse estrogen receptor gene.** *Proc Natl Acad Sci USA* 1993, **90**:11162–11166.

This is the first report of the generation of α ERKO mice and presents a basic description of the phenotypes.

5. Couse JF, Korach KS: **Estrogen receptor null mice: what have we learned and where will they lead us?** *Endocr Rev* 1999, **20**:358–417.

This is a very thorough review of the roles of estrogen in many processes as revealed by the experimental literature as well as data from transgenic mice.

6. Richards JS: **Estradiol receptor content in rat granulosa cells during follicular development: modification by estradiol and gonadotropins.** *Endocrinology* 1975, **97**:1174–1184.

7. Richards JS: **Maturation of ovarian follicles: actions and interactions of pituitary and ovarian hormones on follicular cell differentiation.** *Physiol Rev* 1980, **60**:51–89.

8. Rao MC, Midgley AR Jr, Richards JS: **Hormonal regulation of ovarian cellular proliferation.** *Cell* 1978, **14**:71–78.

9. Goldenberg RL, Vaitukaitis JL, Ross GT: **Estrogen and follicle stimulation hormone interactions on follicle growth in rats.** *Endocrinology* 1972, **90**:1492–1498.

10. Bley MA, Saragueta PE, Baranao JL: **Concerted stimulation of rat granulosa cell deoxyribonucleic acid synthesis by sex steroids and follicle-stimulating hormone.** *J Steroid Biochem Mol Biol* 1997, **62**:11–19.

11. Reilly CM, Cannady WE, Mahesh VB, Stopper VS, De Sevilla LM, Mills TM: **Duration of estrogen exposure prior to follicle-stimulating hormone stimulation is critical to granulosa cell growth and differentiation in rats.** *Biol Reprod* 1996, **54**:1336–1342.

12. Burghardt RC, Anderson E: **Hormonal modulation of gap junctions in rat ovarian follicles.** *Cell Tissue Res* 1981, **214**:181–193.

13. Hsueh AJ, Billig H, Tsafirri A: **Ovarian follicle atresia: a hormonally controlled apoptotic process.** *Endocr Rev* 1994, **15**:707–724.

14. Tonetta SA, Spicer LJ, Ireland JJ: **Cl628 inhibits follicle-stimulating hormone (FSH)-induced increases in FSH receptors of the rat ovary: requirement of estradiol for FSH action.** *Endocrinology* 1985, **116**:715–722.

15. Tonetta SA, diZerega GS: **Intragonadal regulation of follicular maturation.** *Endocr Rev* 1989, **10**:205–229.

16. Kessel B, Liu YX, Jia XC, Hsueh AJ: **Autocrine role of estrogens in the augmentation of luteinizing hormone receptor formation in cultured rat granulosa cells.** *Biol Reprod* 1985, **32**:1038–1050.

17. Wang XN, Greenwald GS: **Synergistic effects of steroids with FSH on folliculogenesis, steroidogenesis and FSH- and hCG-receptors in hypophysectomized mice.** *J Reprod Fertil* 1993, **99**:403–413.

18. Zhuang LZ, Adashi EY, Hsueh AJ: **Direct enhancement of gonadotropin-stimulated ovarian estrogen biosynthesis by estrogen and clomiphene citrate.** *Endocrinology* 1982, **110**:2219–2221.

19. Sar M, Welsch F: **Differential expression of estrogen receptor-beta and estrogen receptor-alpha in the rat ovary.** *Endocrinology* 1999, **140**:963–971.

This study presents a very nice immunohistochemical analysis of localization of the two ERs.

20. Imagawa W, Yang J, Guzman R, Nandi S: **Control of mammary development.** In *The Physiology of Reproduction*. Edited by Knobil E, Neil JD, Ewing LL, Greenwald GS, Markert CL, Pfaff DW. New York: Raven Press, 1994:1033–1063.
21. Lydon JP, DeMayo FJ, Funk CR, Mani SK, Hughes AR, Montgomery CA, Shyamala G, Conneely OM, O'Malley BW: **Mice lacking progesterone receptor exhibit pleiotropic reproductive abnormalities.** *Genes Dev* 1995, **9**:2266–2278.
This is the first report of the generation of PR knockout animals and an initial description of the resulting phenotypes.
22. Korach KS, Couse JF, Curtis SW, Washburn TF, Lindzey J, Kimbro KS, Eddy EM, Migiaccio S, Snedeker SM, Lubahn DB, Schomberg DW, Smith EP: **Estrogen receptor gene disruption: molecular characterization and experimental and clinical phenotypes.** In *Recent Progress in Hormone Research*. Edited by Conn PM. Bethesda, MD: The Endocrine Society, 1996:159–188.
23. Bocchinfuso WP, Korach KS: **Mammary gland development and tumorigenesis in estrogen receptor knockout mice.** *J Mammary Gland Biol Neoplasia* 1997, **2**:323–334.
24. Komm BS, Bodine PV: **The ongoing saga of osteoporosis treatment.** *J Cell Biochem Suppl* 1998, **31**:277–283.
25. Vidal O, Lindberg M, Savendahl L, Lubahn DB, Ritzen EM, Gustafsson JA, Ohlsson C: **Disproportional body growth in female estrogen receptor-alpha-inactivated mice.** *Biochem Biophys Res Commun* 1999, **265**:569–571.
26. Smith EP, Boyd J, Frank GR, Takahashi H, Cohen RM, Specker B, Williams TC, Lubahn DB, Korach KS: **Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man.** *N Engl J Med* 1994, **331**:1056–1061.
27. Barrett-Connor E, Wenger NK, Grady D, Mosca L, Collins P, Kornitzer M, Cox DA, Moscarelli E, Anderson PW: **Hormone and nonhormone therapy for the maintenance of postmenopausal health: the need for randomized controlled trials of estrogen and raloxifene.** *J Womens Health* 1998, **7**:839–847.
28. Karas RH, Baur WE, van Eickles M, Mendelsohn ME: **Human vascular smooth muscle cells express an estrogen receptor isoform.** *FEBS Lett* 1995, **377**:103–108.
29. Johnson BD, Zheng W, Korach KS, Scheuer T, Catterall WA, Rubanyi GM: **Increased expression of the cardiac L-type calcium channel in estrogen receptor-deficient mice.** *J Gen Physiol* 1997, **110**:135–140.
This study shows that the mechanisms of estrogen effects in the heart are ER α mediated.
30. Rubanyi GM, Freay AD, Kauser K, Sukovich D, Burton G, Lubahn DB, Couse JF, Curtis SW, Korach KS: **Vascular estrogen receptors and endothelium-derived nitric oxide production in the mouse aorta. Gender difference and effect of estrogen receptor gene disruption.** *J Clin Invest* 1997, **99**:2429–2437.
31. Iafrazi MD, Karas RH, Aronovitz M, Kim S, Sullivan TR Jr, Lubahn DB, O'Donnell TF Jr, Korach KS, Mendelsohn ME: **Estrogen inhibits the vascular injury response in estrogen receptor alpha-deficient mice.** *Nat Med* 1997, **3**:545–548.
See [32*]
32. Karas RH, Hodgins JB, Kwoun M, Kregge JH, Aronovitz M, Mackey W, Gustafsson JA, Korach KS, Smithies O, Mendelsohn ME: **Estrogen inhibits the vascular injury response in estrogen receptor beta-deficient female mice.** *Proc Natl Acad Sci USA* 1999, **96**:15133–15136.
These two studies [31*,32*], interestingly, indicate that the protective effect of estrogen is not ER mediated, or that either receptor will suffice in the absence of the other.
33. Graham JD, Clarke CL: **Physiological action of progesterone in target tissues.** *Endocr Rev* 1997, **18**:502–519.
34. Curtis SW, Clark J, Myers P, Korach KS: **Disruption of estrogen signaling does not prevent progesterone action in the estrogen receptor or knockout mouse uterus.** *Proc Natl Acad Sci USA* 1999, **96**:3646–3651.
This study indicates that progesterone responses are not dependent on estrogen action and can still occur in the absence of ER α .
35. Cenni B, Picard D: **Ligand-independent activation of steroid receptors: new roles for old players.** *Trends Endocrinol Metab* 1999, **10**:41–46.
This study is a good review of the topic.
36. Curtis SW, Washburn T, Sewall C, DiAugustine R, Lindzey J, Couse JF, Korach KS: **Physiological coupling of growth factor and steroid receptor signaling pathways: estrogen receptor knockout mice lack estrogen-like response to epidermal growth factor.** *Proc Natl Acad Sci USA* 1996, **93**:12626–12630.
This study presents the first demonstration of ligand-independent effects with an *in vivo* system.
37. Nelson KG, Takahashi T, Bossert NL, Walmer DK, McLachlan JA: **Epidermal growth factor replaces estrogen in the stimulation of female genital-tract growth and differentiation.** *Proc Natl Acad Sci USA* 1991, **88**:21–25.
38. Ignar Trowbridge DM, Nelson KG, Bidwell MC, Curtis SW, Washburn TF, McLachlan JA, Korach KS: **Coupling of dual signaling pathways: epidermal growth factor action involves the estrogen receptor.** *Proc Natl Acad Sci USA* 1992, **89**:4658–4662.
This study demonstrates that, biochemically, the ER α responds in a similar manner to EGF and estrogen.
39. Ignar Trowbridge DM, Teng CT, Ross KA, Parker MG, Korach KS, McLachlan JA: **Peptide growth factors elicit estrogen receptor-dependent transcriptional activation of an estrogen-responsive element.** *Mol Endocrinol* 1993, **7**:992–998.
40. Ignar-Trowbridge DM, Pimentel M, Teng CT, Korach KS, McLachlan JA: **Cross talk between peptide growth factor and estrogen receptor signaling systems.** *Environ Health Perspect* 1995, **103** (suppl 7): 35–38.
41. Ignar-Trowbridge DM, Pimentel M, Parker MG, McLachlan JA, Korach KS: **Peptide growth factor cross-talk with the estrogen receptor requires the A/B domain and occurs independently of protein kinase C or estradiol.** *Endocrinology* 1996, **137**:1735–1744.
42. Couse JF, Hewitt SC, Bunch DO, Sar M, Walker VR, Davis BJ, Korach KS: **Postnatal sex reversal of the ovaries in mice lacking estrogen receptors alpha and beta.** *Science* 1999, **286**:2328–2331.
This study is the first report of generation of $\alpha\beta$ ERKO mice and presents a description of their unique ovarian phenotype.
43. Couse JF, Curtis SW, Washburn TF, Lindzey J, Golding TS, Lubahn DB, Smithies O, Korach KS: **Analysis of transcription and estrogen insensitivity in the female mouse after targeted disruption of the estrogen receptor gene.** *Mol Endocrinol* 1995, **9**:1441–1454.
This study provides a further description of the estrogen insensitivity of the α ERKO mouse.

Authors' affiliation: Receptor Biology, LRDT, NIEHS, National Institutes of Health, Research Triangle Park, North Carolina, USA

Correspondence: Kenneth S Korach, Receptor Biology, LRDT, NIEHS, National Institutes of Health, Research Triangle Park, NC 27709, USA. Tel: +1 919 541 312; fax: +1 919 541 0696; e-mail: korach@niehs.nih.gov